

Dirac Equation in Noncommutative Space for Hydrogen Atom

T. C. Adorno¹, M. C. Baldiotti¹, M. Chaichian²,
D. M. Gitman¹ and A. Tureanu²

¹*Instituto de Física, Universidade de São Paulo, Caixa Postal 66318,
CEP 05508-090 São Paulo, S.P., Brazil*

²*Department of Physics, University of Helsinki and Helsinki Institute of Physics,
P.O.Box 64, FIN-00014 Helsinki, Finland*

Abstract

We consider the energy levels of a hydrogen-like atom in the framework of θ -modified, due to space noncommutativity, Dirac equation with Coulomb field. It is shown that on the noncommutative (NC) space the degeneracy of the levels $2S_{1/2}$, $2P_{1/2}$ and $2P_{3/2}$ is lifted completely, such that new transition channels are allowed.

1 Introduction

Possible physically observable consequences of noncommutativity of space coordinates have attracted recently a lot of attention in order to justify the intensive study of noncommutative versions of QFT and quantum mechanics, mainly motivated by the seminal works [1, 2]. The results of Ref. [3] show, e.g., that the noncommutativity leads to a deviation of the energy levels of the hydrogen atom from the well-known structure obtained in the framework of standard quantum mechanics. The results of [3] show that the degenerate levels $2S - 2P$ split in three others levels. However, the calculations of [3] were done in the framework of θ -modified nonrelativistic Schrödinger equation with Coulomb field. It is interesting to refine these results, considering a noncommutative modification of the energy levels of the hydrogen atom in the framework of the θ -modified Dirac equation with the Coulomb field, which is the relativistic Schrödinger equation for the case under consideration. Such a problem is

solved in the present work. We show that in contrast with the nonrelativistic case, the noncommutativity lifts completely the degeneracy of the levels $2P_{1/2}$ and $2P_{3/2}$, opening new allowed transition channels.

2 θ -modified Dirac equation with Coulomb field

The behavior of an electron with charge $-e$ ($e > 0$) and mass m in the Coulomb field of a nucleus Ze , in a noncommutative space, is determined by the θ -modified Dirac equation, whose Hamiltonian form is

$$\begin{aligned} i\frac{\partial}{\partial t}\Psi(x) &= \hat{H}^\theta\Psi(x), \quad \hat{H}^\theta = (\boldsymbol{\alpha} \cdot \hat{\mathbf{p}}) + m\gamma^0 - eA_0(q), \\ \alpha_i &\equiv \gamma_0\gamma_i, \quad \boldsymbol{\alpha} = (\alpha_i), \quad \hat{\mathbf{p}} = (p_i), \end{aligned} \quad (1)$$

where $\Psi(x)$ is a four component spinor, $\hat{\mathbf{p}} = -i\nabla$, γ_μ are the Dirac matrices and the θ -modified Coulomb potential $A_0(q)$ reads

$$A_0(q) = A_0\left(x^i - \frac{1}{2}\theta^{ij}\hat{p}_j\right) = \frac{Ze}{\sqrt{\left(x^i - \frac{1}{2}\theta^{ij}\hat{p}_j\right)\left(x^i - \frac{1}{2}\theta^{ik}\hat{p}_k\right)}}, \quad i, j, k = (1, 2, 3), \quad (2)$$

where θ^{jk} is the constant anti-symmetric noncommutativity parameter that defines the algebra of noncommutative position operators q^i , $[q^j, q^k] = i\theta^{jk}$ (see, e.g., [4]). In what follows we call $\hat{H}^\theta$ the θ -modified Dirac Hamiltonian with Coulomb field on a noncommutative space. Setting $\theta^i = \varepsilon_{ijk}\theta^{jk}$ (ε_{ijk} is the Levi-Civita symbol with the usual definition) and denoting $r = \sqrt{x^i x^i}$, the potential (2) can be written as

$$A_0\left(x^i - \frac{1}{2}\theta^{ij}\hat{p}_j\right) = \frac{Ze}{r} + \frac{1}{4}\frac{Ze}{r^3}(\hat{\mathbf{L}} \cdot \boldsymbol{\theta}) + O(\theta^2), \quad \boldsymbol{\theta} = (\theta^i),$$

where $\hat{\mathbf{L}} = [\mathbf{x} \times \hat{\mathbf{p}}]$ is the angular momentum operator. Therefore the θ -modified Dirac Hamiltonian takes the form

$$\hat{H}^\theta = \hat{H} + \hat{V}^\theta + O(\theta^2), \quad (3)$$

$$\hat{H} = (\boldsymbol{\alpha} \cdot \mathbf{p}) + m\gamma^0 - \frac{Ze^2}{r}, \quad \hat{V}^\theta = -\frac{1}{4}\frac{Ze^2}{r^3}(\hat{\mathbf{L}} \cdot \boldsymbol{\theta}). \quad (4)$$

Using the exact eigenfunctions of $\hat{H}$ and treating $\hat{V}^\theta$ as a perturbation of the ordinary Dirac Hamiltonian $\hat{H}$, one can calculate the modification of energy levels of the hydrogen atom in the framework of θ -modified Dirac equation. The spectrum of $\hat{H}$ and the corresponding eigenfunctions are well known, see [5, 6, 7, 9]. Below, we list the spectrum and the

eigenfunctions of $\hat{H}$ for $Z < 137$:

$$\begin{aligned}\hat{H}\Psi_{N,j,M,\zeta} &= E_{N,j}\Psi_{N,j,M,\zeta}, \quad \zeta = \pm 1, \quad E_{N,j} = \frac{m}{\sqrt{1 + \left(\frac{Ze^2}{\gamma+N}\right)^2}}, \\ N &= \begin{cases} 1, 2, \dots, & \zeta = 1 \\ 0, 1, 2, \dots, & \zeta = -1 \end{cases}, \quad \gamma = \sqrt{\kappa^2 - (Ze^2)^2}, \quad \kappa = \zeta \left(j + \frac{1}{2}\right), \\ \hat{\mathbf{J}}^2\Psi_{N,j,M,\zeta} &= j(j+1)\Psi_{N,j,M,\zeta}, \quad \hat{J}_3\Psi_{N,j,M,\zeta} = M\Psi_{N,j,M,\zeta}, \\ \hat{\mathbf{J}} &= \hat{\mathbf{L}} + \frac{1}{2}\boldsymbol{\Sigma}, \quad \boldsymbol{\Sigma} = \text{diag}(\boldsymbol{\sigma}, \boldsymbol{\sigma}), \quad j = \frac{1}{2}, \frac{3}{2}, \dots, \quad -j \leq M \leq j,\end{aligned}$$

where $\boldsymbol{\sigma} = (\sigma_i)$ are the Pauli matrices and the bispinors $\Psi_{N,j,M,\zeta}$ are given by

$$\begin{aligned}\Psi_{N,j,M,\zeta}(r, \vartheta, \varphi) &= \frac{1}{r} \begin{pmatrix} \Omega_{j,M,\zeta}(\vartheta, \varphi) F_{N,j,\zeta}^+(r) \\ i\Omega_{j,M,-\zeta}(\vartheta, \varphi) F_{N,j,\zeta}^-(r) \end{pmatrix}, \\ \Omega_{j,M,\zeta}(\vartheta, \varphi) &= \frac{\zeta}{\sqrt{2j+\zeta+1}} \begin{pmatrix} -\sqrt{(j+\frac{\zeta}{2}) - \zeta M + \frac{1}{2}} Y_{j+\frac{\zeta}{2}, M-\frac{1}{2}}(\vartheta, \varphi) \\ \sqrt{(j+\frac{\zeta}{2}) + \zeta M + \frac{1}{2}} Y_{j+\frac{\zeta}{2}, M+\frac{1}{2}}(\vartheta, \varphi) \end{pmatrix},\end{aligned}\tag{5}$$

where $Y_{A,B}(\vartheta, \varphi)$ are the spherical harmonics [6, 7] and

$$\begin{aligned}F_{N,j,\zeta}^\pm(r) &= \pm A \lambda_N^\pm z^{\gamma-1} e^{-z/2} [(\eta_N - \kappa) \Phi(-N, \beta; z) \mp N \Phi(-N+1, \beta; z)], \\ A &= \frac{(2\lambda_N)^{3/2}}{2\Gamma(\beta)} \sqrt{\frac{\Gamma(\beta+N)}{\eta_N(\eta_N - \kappa)N!}}, \quad \lambda_N^\pm = \sqrt{1 \pm \frac{E_{N,j}}{m}}, \quad \lambda_N = m\lambda_N^+ \lambda_N^-, \\ \beta &= 2\gamma + 1, \quad \eta_N = Ze^2 m / \lambda_N, \quad z = 2\lambda_N r,\end{aligned}\tag{6}$$

where $\Phi(a, b; z)$ is the hypergeometric confluent function (see [10] for definition).

3 θ -modification of energy levels

As it was shown in [3], for the hydrogenic atoms, without spin-orbit interaction, the nonrelativistic energy levels of the θ -modified Hamiltonian $\hat{H}^{\text{nr}}$,

$$\hat{H}^{\text{nr}} = \frac{\hat{\mathbf{p}}^2}{2m} - \frac{Ze^2}{r},\tag{7}$$

are characterized by a lift of degeneracy in the quantum number $l = j + \zeta/2$ (eigenvalue of $\hat{\mathbf{L}}^2$ operator), implying certain splittings of the energy levels. Thus, there appears a possibility of new transition channels with distinct projections of total angular momentum M , i.e., in the spectroscopic notation, transitions like $nl_j^M \longrightarrow nl_j^{M'}$ (for $l \neq 0$), where $n = N + |\kappa|$ is the nonrelativistic principal quantum number, eigenvalue of the radial differential operator

of (7). The perturbation due to the noncommutativity is given by the same operator $\hat{V}^\theta$ as in (4), but it is calculated with the nonrelativistic wave functions, whose correction on degenerate levels, in leading order, is obtained by computing the eigenvalues of the secular matrix ΔE^{nr} , its elements being defined as

$$\Delta E_{aa'}^{\text{nr}} = -\frac{Ze^2}{4} \int_0^\infty dr r^2 \int_0^{4\pi} d\Omega \left\{ \psi_{n,a}^\dagger(r, \vartheta, \varphi) \left[\frac{\hat{\mathbf{L}} \cdot \boldsymbol{\theta}}{r^3} \right] \psi_{n,a'}(r, \vartheta, \varphi) \right\}, \quad (8)$$

where the label $a = (l, j, M)$ is the collection of the three degenerated nonrelativistic quantum numbers and

$$\begin{aligned} \psi_{n,a}(r, \vartheta, \varphi) &\equiv \psi_{n,l,j,M}(r, \vartheta, \varphi) = R_{n,l}(r) \chi_{j,M}(\vartheta, \varphi), \\ \hat{j}^2 \chi_{j,M}(\vartheta, \varphi) &= \left(\hat{\mathbf{L}} + \boldsymbol{\sigma}/2 \right)^2 \chi_{j,M}(\vartheta, \varphi) = j(j+1) \chi_{j,M}(\vartheta, \varphi) \end{aligned}$$

are the eigenfunctions of (7), $\hat{H}^{\text{nr}} \psi_{n,a} = E_n \psi_{n,a}$.

In order to obtain the θ -modification of the $2S_{1/2}$, $2P_{1/2}$, $2P_{3/2}$ levels, we emphasize that $\hat{V}^\theta$ is a vectorial operator and, according to angular momentum selections rules, allowed transitions are between levels with $\Delta M \equiv |M - M'| = 0, 1$ and $\Delta l \equiv |l - l'| = 0$ [8]. The 6×6 secular matrix ΔE^{nr} has three degenerate eigenvalues represented as ΔE_0^{nr} and $\Delta E_\pm^{\text{nr}}$ and, restoring $\hbar$ and c , these corrections have the form

$$\Delta E_0^{\text{nr}} = 0, \quad \Delta E_\pm^{\text{nr}} = -\frac{mc^2}{4} \left(\frac{Z^2 \alpha^2}{\lambda_e} \right)^2 \left(\frac{1}{24} \right) (\pm |\boldsymbol{\theta}|), \quad |\boldsymbol{\theta}| = \sqrt{\theta^i \theta^i}, \quad (9)$$

where $\alpha = e^2/\hbar c$ is the fine structure constant, $\lambda_e = \lambda_e/2\pi = \hbar/mc$ and λ_e is the electron Compton wavelength. The same results are obtained if we choose $\theta_1 = \theta_2 = 0$ and $\theta_3 \neq 0$ (which can be done by a rotation or a redefinition of coordinates). With this choice, $|\boldsymbol{\theta}| = \theta_3$, and the diagonal elements can be calculated using the general formulae obtained in [3]. Since the goal of [3] is the study of the transition $^1 2P_{1/2} \rightarrow 2S_{1/2}$, it does not contain the expression for the nondiagonal elements of $\Delta E_{bb'}^{\text{nr}}$. For these nondiagonal elements, we have

$$\begin{aligned} \Delta E_{bb'}^{\text{nr}} &= -\frac{Ze^2}{4} \int_0^\infty dr r^2 \int_0^{4\pi} d\Omega \left\{ \psi_{2,b}^\dagger(r, \vartheta, \varphi) \left[\frac{\hat{\mathbf{L}} \cdot \boldsymbol{\theta}}{r^3} \right] \psi_{2,b'}(r, \vartheta, \varphi) \right\} \\ &= -\frac{mc^2}{4} \left(\frac{Z^2 \alpha^2}{\lambda_e} \right)^2 \left(\frac{1}{24} \right) \left(\frac{\sqrt{2}}{3} \theta_3 \right), \quad b = \left(1, \frac{1}{2}, \pm \frac{1}{2} \right), \quad b' = \left(1, \frac{3}{2}, \pm \frac{1}{2} \right). \end{aligned} \quad (10)$$

¹Rigorously this transition does not exist, as the perturbed level is actually a superposition of the degenerated levels $2P_{1/2}$ and $2P_{3/2}$, and only the quantum number M is well defined. But this consideration does not change the numerical value of the transition and, consequently, could be ignored in [3].

With this definition of coordinates, $[\hat{J}_3, \hat{V}^\theta] = 0$, therefore the eigenstates of $\hat{H}^{\theta, \text{nr}} = \hat{H}^{\text{nr}} + \hat{V}^\theta$, in leading order, have the magnetic quantum number M well defined. This result allows us to conclude that, in nonrelativistic case, the $2S - 2P$ levels split in three other levels, all of them *twofold degenerate*, as illustrated in Figure 1.

In the relativistic theory, the θ -modification of the energy levels are obtained by computing the eigenvalues of the secular matrix ΔE^{rel} , characterized by the mean values of the operator $\hat{V}^\theta$ (4), but now with respect to the Dirac spinors $\Psi_{N,j,M,\zeta}$ (5), with the same angular momentum selection rules, i.e., $\Delta M \equiv |M - M'| = 0, 1$ and $\Delta l \equiv |l - l'| = 0$. The perturbation operator $\hat{V}^\theta$ does not mix the small $F_{N,j,\zeta}^-(r)$ and the big $F_{N,j,\zeta}^+(r)$ components of the Dirac spinor (5), so ΔE^{rel} consists of two terms, which read:

$$\begin{aligned} \Delta E_{\mu\mu'}^{\text{rel}} = & -\frac{Ze^2}{4} \int_0^\infty \frac{dr}{r} [(F_{N,j,\zeta}^+)^* F_{N,j,\zeta'}^+] \int_0^{4\pi} d\Omega \left\{ \Omega_{j,M,\zeta}^\dagger (\hat{\mathbf{L}} \cdot \boldsymbol{\theta}) \Omega_{j,M',\zeta'} \right\} \\ & - \frac{Ze^2}{4} \int_0^\infty \frac{dr}{r} [(F_{N,j,\zeta}^-)^* F_{N,j,\zeta'}^-] \int_0^{4\pi} d\Omega \left\{ \Omega_{j,M,-\zeta}^\dagger (\hat{\mathbf{L}} \cdot \boldsymbol{\theta}) \Omega_{j,M',-\zeta'} \right\}, \end{aligned}$$

where the label $\mu = (M, \zeta)$ is the set of two degenerated relativistic quantum numbers. However, $F_{N,j,\zeta}^-(r)$ is approximately by $1/c$ smaller than $F_{N,j,\zeta}^+(r)$, so we can neglect the terms proportional to small components, retaining only the first term above.

We are interested in calculating the θ -modifications of the relativistic energy of the $2S, 2P$ levels. The nonrelativistic degeneracy in the quantum number j is naturally removed in the Dirac theory, in such a way that the level $2P_{3/2}$ is lifted from the still degenerate $2S_{1/2}, 2P_{1/2}$ levels. The space noncommutativity leads to additional splittings and a separate analysis for the $j = 1/2$ and $j = 3/2$ relativistic energy levels is required. The elements of the secular matrix $\Delta E^{\text{rel}}(j)$, in leading order, are

$$\Delta E_{MM'}^{\text{rel}}(j) = -\frac{Ze^2}{4} \int_0^\infty dr r^2 |F_{N,j,\zeta}^+|^2 \int_0^{4\pi} d\Omega \left\{ \Omega_{j,M,\zeta}^\dagger \left[\frac{\hat{\mathbf{L}} \cdot \boldsymbol{\theta}}{r^3} \right] \Omega_{j,M',\zeta} \right\}, \quad (11)$$

whose results, for each level, follow below.

3.1 Relativistic NC correction for the $2P_{1/2}$ level

The θ -correction for the $2P_{1/2}$ level ($N = 1, j = 1/2, \zeta = 1, M = \pm \frac{1}{2}$) follows from (11):

$$\begin{aligned} \Delta E_{MM'}^{\text{rel}}(1/2) = & -\frac{Ze^2}{4} \varrho_{1/2} \Theta_{MM'}^{\text{rel}}, \\ \varrho_{1/2} = & \int_0^\infty \frac{dr}{r} \left| F_{1,\frac{1}{2},1}^+(r) \right|^2, \quad \Theta_{MM'}^{\text{rel}} = \int_0^{4\pi} d\Omega \left[\Omega_{\frac{1}{2},M,1}^* (\hat{\mathbf{L}} \cdot \boldsymbol{\theta}) \Omega_{\frac{1}{2},M',1} \right]. \end{aligned} \quad (12)$$

Expressing the hypergeometric confluent functions (6) in terms of exponentials and polynomials, we obtain

$$\begin{aligned} \varrho_{1/2} &= (2\lambda_1)^3 \left[\frac{\beta_1 (\lambda_1^+)^2}{4\eta_1 (\eta_1 - 1) (\beta_1 - 1) (\beta_1 - 2) (\beta_1 - 3)} \right] \{ (\eta_1 - 2)^2 \\ &\quad - \left[\frac{2(\eta_1 - 2)(\eta_1 - 1)(\beta_1 - 3)}{\beta_1} \right] + \left(\frac{\eta_1 - 1}{\beta_1} \right)^2 (\beta_1 - 2)(\beta_1 - 3) \} , \\ \eta_1 &= Ze^2 \frac{m}{\lambda_1}, \quad \gamma_1 = \sqrt{1 - (Ze^2)^2}, \quad \beta_1 = 2\gamma_1 + 1, \end{aligned} \quad (13)$$

with λ_1^+ , λ_1 given in (6). The matrix Θ^{rel} , whose elements are $\Theta_{MM'}^{\text{rel}}$, reads

$$\Theta^{\text{rel}} = \frac{2}{3} \begin{pmatrix} -\theta_3 & \theta_+ \\ \theta_- & \theta_3 \end{pmatrix}, \quad \theta_{\pm} = \theta_1 \pm i\theta_2, \quad (14)$$

and its eigenvalues, denoted by $\Lambda_j^{\pm}$, are

$$\Lambda_{1/2}^{\pm} = \pm \frac{2}{3} |\boldsymbol{\theta}|, \quad |\boldsymbol{\theta}| = \sqrt{\theta^i \theta^i}. \quad (15)$$

The θ -corrections to the relativistic energy of the $2P_{1/2}$ level, $\Delta E_{\pm}^{\text{rel}}(j)$, restoring $\hbar$ and c , reads

$$\begin{aligned} \Delta E_{\pm}^{\text{rel}}(1/2) &= -\frac{Ze^2}{4} \varrho_{1/2} \Lambda_{1/2}^{\pm} \\ &= -mc^2 \left(\frac{Z^2 \alpha^2}{\lambda_e} \right)^2 \left(\pm \frac{|\boldsymbol{\theta}|}{3} \right) \left\{ \left(\frac{(\lambda_1^+)^2 \beta_1}{(\beta_1 - 1)(\beta_1 - 2)(\beta_1 - 3)} \right) \right. \\ &\quad \times \left(\frac{1}{\eta_1^4 (\eta_1 - 1)} \right) \left[(\eta_1 - 2)^2 - \left(\frac{2(\eta_1 - 2)(\eta_1 - 1)(\beta_1 - 3)}{\beta_1} \right) \right. \\ &\quad \left. \left. + \left(\frac{\eta_1 - 1}{\beta_1} \right)^2 (\beta_1 - 2)(\beta_1 - 3) \right] \right\}. \end{aligned} \quad (16)$$

Therefore there is a splitting of this level in such way that the originally degenerate level $2S_{1/2}$, $2P_{1/2}$ splits in three sublevels. Similarly to the nonrelativistic case, the same results can be obtained by choosing a reference frame where $\theta_1 = \theta_2 = 0$ and $\theta_3 \neq 0$. The advantage of this frame is that $[\hat{J}_3, \hat{V}^{\theta}] = 0$ and therefore, the new states have the magnetic quantum number M well defined. In this reference frame, the splitting of the level $2S_{1/2}$, $2P_{1/2}$ in the three levels $2P_{1/2}^{+1/2}$, $2S_{1/2}$, $2P_{1/2}^{-1/2}$ is illustrated in Figure 1.

3.2 Relativistic NC correction for the $2P_{3/2}$ level

According to (11), the θ -correction for the $2P_{3/2}$ level ($N = 0$, $j = 3/2$, $\zeta = -1$, $M = \pm 1/2, \pm 3/2$) is

$$\Delta E_{MM'}^{\text{rel}}(3/2) = -\frac{Ze^2}{4} \varrho_{3/2} \Theta_{MM'}^{\text{rel}}, \quad (17)$$

where the radial integral is

$$\varrho_{3/2} = \int_0^\infty \frac{dr}{r} \left| F_{0, \frac{3}{2}, -1}^+(r) \right|^2 = 4(\lambda_0)^3 \left[\frac{(\lambda_0^+)^2}{(\beta_2 - 1)(\beta_2 - 2)(\beta_2 - 3)} \right], \quad (18)$$

$$\gamma_2 = \sqrt{4 - (Ze^2)^2}, \quad \beta_2 = 2\gamma_2 + 1,$$

with λ_0^+ , λ_0 given in (6). Compared to the previous case, the matrix Θ^{rel} is now a 4×4 -matrix, whose elements are

$$\Theta_{MM'}^{\text{rel}} = \int_0^{4\pi} d\Omega \left[\Omega_{\frac{3}{2}, M, -1}^* \left(\hat{\mathbf{L}} \cdot \boldsymbol{\theta} \right) \Omega_{\frac{3}{2}, M', -1} \right], \quad (19)$$

and explicitly is given by

$$\Theta^{\text{rel}} = \begin{pmatrix} -\theta_3 & \frac{1}{\sqrt{3}}\theta_+ & 0 & 0 \\ \frac{1}{\sqrt{3}}\theta_- & -\frac{1}{3}\theta_3 & \frac{2}{3}\theta_+ & 0 \\ 0 & \frac{2}{3}\theta_- & \frac{1}{3}\theta_3 & \frac{1}{\sqrt{3}}\theta_+ \\ 0 & 0 & \frac{1}{\sqrt{3}}\theta_- & \theta_3 \end{pmatrix}, \quad \theta_\pm = \theta_1 \pm i\theta_2. \quad (20)$$

Its four nondegenerate eigenvalues, $\Lambda_j^\pm$ and $\Lambda_j^{\pm'}$, are

$$\Lambda_{3/2}^\pm = \pm |\boldsymbol{\theta}|, \quad \Lambda_{3/2}^{\pm'} = \frac{\Lambda_{3/2}^\pm}{3}, \quad |\boldsymbol{\theta}| = \sqrt{\theta^i \theta^i}. \quad (21)$$

The θ -corrections for the $2P_{3/2}$ level, in dimension of energy (restoring $\hbar$ and c), have the form

$$\Delta E^{\text{rel}}(3/2)^\pm = -\frac{Ze^2}{4} \varrho_{3/2} \Lambda_{3/2}^\pm, \quad \Delta E^{\text{rel}}(3/2)^{\pm'} = \frac{\Delta E^{\text{rel}}(3/2)^\pm}{3},$$

$$\Delta E^{\text{rel}}(3/2)^\pm = -\frac{mc^2}{4} \left(\frac{Z^2 \alpha^2}{\lambda_e} \right)^2 (\pm |\boldsymbol{\theta}|) \frac{(\lambda_0^+)^2}{2(\beta_2 - 1)(\beta_2 - 2)(\beta_2 - 3)}. \quad (22)$$

Choosing again the reference frame where $\theta_1 = \theta_2 = 0$ and $\theta_3 \neq 0$, we have $[\hat{J}_3, \hat{V}^\theta] = 0$ and the level $2P_{3/2}$ splits in the four levels $2P_{3/2}^{+1/2}$, $2P_{3/2}^{-1/2}$, $2P_{3/2}^{+3/2}$, $2P_{3/2}^{-3/2}$, as illustrated in Figure 1.

The energy levels and their modifications due to presence of space noncommutativity, for the relativistic and nonrelativistic cases, are shown² in Figure 1.

²On this figure, the energy splits indicated by the dashed-point arrow comes from the usual Dirac equation, so it does not depends on θ and this split can not be compared with the others θ -modified levels.




Figure 1 – Splittings for relativistic and nonrelativistic energy levels due to space noncommutativity. All numerical values of the coefficients of the first-order θ -corrections are in units of eV/m^2 .

We conclude that, from the point of view of the θ -modified relativistic Dirac theory there is an additional, in contrast to the nonrelativistic case, splitting of some degenerate levels and there appear new transition channels. In particular, in the presence of the space noncommutativity, the degenerate levels $2S_{1/2}, 2P_{1/2}$ split into three nondegenerate sublevels and the level $2P_{3/2}$ splits into four nondegenerate sublevels, and the transition $2P_{1/2}^{\pm 1/2} \rightarrow 2S_{1/2}$ is possible. Except for the spherically symmetric levels $2S_{1/2}^{\pm 1/2}$, these results show that in the noncommutative relativistic theory degeneracy is completely removed. The obtained results show explicitly the dependence of the energy levels of the hydrogen atom on the noncommutative parameter. Once these energy levels can be measured experimentally with a high accuracy, the actual spectroscopy data can be used to impose bounds on the value of the noncommutativity parameter θ .

Acknowledgements

T.C.A. thanks CNPq; M.C.B. thanks FAPESP; D.M.G. thanks FAPESP and CNPq for permanent support. The support of the Academy of Finland under the Projects no. 121720 and 127626 is greatly acknowledged.

References

- [1] S. Doplicher, K. Fredenhagen and J. E. Roberts, Phys. Lett. B **331** (1994) 39; Commun. Math. Phys. **172** (1995) 187, hep-th/0303037.
- [2] N. Seiberg and E. Witten, JHEP **09** (1999) 032, hep-th/9908142.
- [3] M. Chaichian, M. M. Sheikh-Jabbari and A. Tureanu, Phys. Rev. Lett. **86** (2001) 2716.
- [4] D. M. Gitman and V. G. Kupriyanov, Eur. Phys. J. C **54** (2008) 325-332.
- [5] H. A. Bethe and E. E. Salpeter, *Quantum Mechanics of One - and Two-Electron Atoms* (Springer - Verlag 1957).
- [6] M. E. Rose, *Relativistic Electron Theory* (John Wiley & Sons 1961).
- [7] A. I. Akhiezer and V. B. Berestetskii, *Quantum Electrodynamics* (Interscience 1965).
- [8] L. D. Landau and E. M. Lifshitz, *Quantum Mechanics* (Pergamon Press 1977).
- [9] B. Voronov, D. M. Gitman and I. V. Tyutin, Teor. Math. Phys. **150** (1) (2007) 34.
- [10] I. S. Gradshteyn and N. M. Ryzhik, *Tables of integrals, sums, series and products* (70th ed. Academic Press 2007).